Applicant: Eleftheria Maraissa Flier et al.

Serial No.: 09/159,068

Filed: September 23, 1998

Page: 2

Add new claims 47-63.

47. A method of inhibiting appetite, or the gain of weight, in a subject comprising: identifying a subject in need of inhibiting appetite or weight gain; and administering an effective amount of an antagonist of melanocyte concentrating hormone (MCH) to said subject, wherein the antagonist binds an MCH receptor.

Attorney's Docker No.: 10276-014002 / JDP-025CN

- 48. The method of claim 47, wherein the antagonist binds competitively to an MCH receptor.
- 49. The method of claim 47, wherein the antagonist binds non-competitively to an MCH receptor.
 - 50. The method of claim 47, wherein the antagonist is a peptide analog of MCH.
- 51. The method of claim 50, wherein the peptide analog has 1, 2, 3, 4, 5 or more residues within the MCH ring structure modified or substituted with a nonconserved amino acid.
- 52. The method of claim 51, wherein between 1 and 5 residues within the MCH ring structure modified or substituted with a nonconserved amino acid.
- 53. The method of claim 50, wherein the peptide analog is deleted for any or all of the residues between R1 and R6 or between R18 and R19.
- 54. The method of claim 51, wherein the peptide analog is deleted for any or all of the residues between R1 and R6 or between R18 and R19.
- 55. The method of claim 52, wherein the peptide analog is deleted for any or all of the residues between R1 and R6 or between R18 and R19.
- 56. The method of claim 50, wherein the peptide analog has least 50 % homology with MCH.
- 57. The method of claim 50, wherein the peptide analog has least 60 % homology with MCH.



Applicant: Eleftheria Maratta Flier et al.

Serial No.: 09/159,068

Filed: September 23, 1998

Page: 3

58. The method of claim 50, wherein the peptide analog has least 70 % homology with MCH.

Attorney's Dock

No.: 10276-014002 / JDP-025CN

59. The method of claim 50, wherein the peptide analog has least 80 % homology with MCH.

60. The method of claim 50, wherein the peptide analog has least 90 % homology with MCH.

61. The method of claim 47, wherein the antagonist is a non-polypeptide drug or chemical.

62. The method of claim 61, wherein the antagonist binds competitively with MCH.

63. The method of claim 61, wherein the antagonist binds non-competitively with MCH.--

B2 Cont